

Depression during pregnancy

Deirdre Ryan, MB, FRCPC Lisa Milis Nicholas Misri

ABSTRACT

OBJECTIVE To review existing literature on depression during pregnancy and to provide information for family physicians in order to promote early detection and treatment.

QUALITY OF EVIDENCE MEDLINE was searched from January 1989 through August 2004 using the key words depression, pregnancy, prenatal, and antenatal. Articles focusing on depression during pregnancy were chosen for review; these articles were based on expert opinion (level III evidence) and prospective studies (level II evidence).

MAIN MESSAGE Pregnancy does not safeguard women against depressive illness. The Edinburgh Postnatal Depression Scale is an effective screening tool for identifying women with depressive symptoms during pregnancy. Once diagnosed with major depression, these patients need to be monitored closely for up to a year after delivery. Patients with mild-to-moderate illness should be referred for psychotherapy. More severely ill patients might require additional treatment with antidepressants. The most commonly used antidepressants are selective serotonin reuptake inhibitors and the serotonin and norepinephrine reuptake inhibitor, venlafaxine. For each patient, risk of treatment with an antidepressant needs to be compared with risk of not treating her depressive illness.

CONCLUSION Early detection of depression during pregnancy is critical because depression can adversely affect birth outcomes and neonatal health and, if left untreated, can persist after the birth. Untreated postpartum depression can impair mother-infant attachments and have cognitive, emotional, and behavioural consequences for children.

RÉSUMÉ

OBJECTIF Recenser les publications traitant de la dépression chez la femme enceinte et donner au médecin de famille l'information nécessaire pour mieux déceler et traiter cette condition.

QUALITÉ DES PREUVES Une recherche a été effectuée dans MEDLINE entre janvier 1989 et août 2004 à l'aide des mots-clés *depression, pregnancy, prenatal, et antenatal*. Les articles centrés sur la dépression durant la grossesse ont été retenus pour analyse; ces articles reposaient sur l'opinion d'experts (preuves de niveau III) et sur des études prospectives (preuves de niveau II).

PRINCIPAL MESSAGE Les femmes enceintes ne sont pas à l'abri des troubles dépressifs. L'échelle de dépression postnatale d'Édimbourg est un outil efficace pour identifier les femmes qui présentent des symptômes dépressifs durant la grossesse. En cas de dépression sévère, la femme devra être suivie de près jusqu'à un an après l'accouchement. Les cas de dépression légère à modérée devraient être dirigés en psychothérapie. Les dépressions plus graves pourraient exiger l'usage d'antidépresseurs. Les plus couramment utilisés sont les inhibiteurs sélectifs du recaptage de la sérotonine et l'inhibiteur du recaptage de la noradrénaline et de la sérotonine, la venlafaxine. Dans chaque cas, le risque lié à l'utilisation de l'antidépresseur doit être comparé à celui de ne pas traiter la dépression.

CONCLUSION Il est crucial de déceler la dépression tôt durant la grossesse; en effet, cette maladie peut affecter négativement l'issue de la grossesse et la santé néonatale; non traitée, elle peut se prolonger au-delà de l'accouchement. Une dépression postnatale peut interférer avec le développement des liens mère-enfant et avoir des conséquences cognitives, émotionnelles et comportementales pour l'enfant.

This article has been peer reviewed
Cet article a fait l'objet d'une révision par des pairs.
Can Fam Physician 2005;51:1087-1093.

There is growing evidence to challenge the notion that pregnancy protects patients from mental illness. Rates of depression during pregnancy have been reported to be as high as 20%.¹ Evidence suggests that postpartum depression (PPD) can be part of a continuum, with onset of illness during pregnancy.¹⁻³ Family physicians are often the first caregivers to come into contact with pregnant women; they, therefore, have a unique opportunity to monitor women's moods over several weeks. Early detection of symptoms could facilitate timely treatment and prevent ongoing depression. This article was written to assist family physicians in recognizing prenatal depression and to aid their decision making for managing this complex condition.

Quality of evidence

MEDLINE was searched from January 1989 through August 2004 using the key words depression, pregnancy, prenatal, and antenatal. Articles were also gathered from the references of papers generated by the initial search. For this article, the search was limited to English-language human studies; articles focusing on depression during pregnancy were chosen for review. The articles were based on expert opinion (level III evidence) and prospective studies (level II evidence). There are few trials offering level I evidence in this area due to ethical restrictions on extracting data on pregnant populations. Available data have been incorporated into this review.

Course of depression during pregnancy

Diagnosis of major depression is often made by mental health professionals during structured clinical interviews using criteria described in

the *Diagnostic and Statistical Manual of Mental Disorders*, fourth edition, text revision (DSM-IV-TR).⁴ For family physicians facing time constraints, one of the quickest ways to assess pregnant women's moods is through responses to self-reported questionnaires in conjunction with objective questions to assess mood.

Screening methods, such as the Edinburgh Postnatal Depression Scale (EPDS) (**Figure 1**),⁵ have been developed for assessing postpartum depression. As yet, no screening tools are specifically designed for assessing antenatal depression. Murray and Cox⁶ investigated use of the EPDS during pregnancy and found that it was effective at identifying women with major depression (level II evidence). The EPDS is commonly used in research to detect perinatal mood disorders.⁷⁻⁹ Because the EPDS is only a screening tool, high scores do not in themselves confirm diagnosis of depression. A score above 12 indicates probable depressive disorder.^{6,7} Other self-rating screening tools, such as the Beck Depression Inventory (BDI),¹⁰ tend to focus on somatic symptoms and, therefore, make detection of depression during pregnancy difficult. We need screening tools that take into account the overlap between symptoms of pregnancy and symptoms of depression.

During the first trimester, it can be particularly difficult to diagnose depression because of this overlap. Bennett et al⁹ reported a rate of depression of 7.4% during the first trimester. During the second trimester, this figure rose to 12.8% and remained at 12% during the third trimester (level I evidence). In a longitudinal study, Josefsson et al⁸ examined rates of depression during late pregnancy and reported

Levels of evidence

Level I: At least one properly conducted randomized controlled trial, systematic review, or meta-analysis

Level II: Other comparison trials, non-randomized, cohort, case-control, or epidemiologic studies, and preferably more than one study

Level III: Expert opinion or consensus statements

Dr Ryan is a Clinical Assistant Professor in the Department of Psychiatry at the University of British Columbia and is a Psychiatrist with the Reproductive Mental Health Program at St Paul's Hospital and BC Women's Hospital and Health Centre in Vancouver.

Ms Milis and Mr Misri are researchers in the Reproductive Mental Health Programs at St Paul's Hospital and BC Women's Hospital and Health Centre.

Figure 1. Edinburgh Postnatal Depression Scale**INSTRUCTIONS FOR USERS**

1. The mother is asked to underline the response that comes closest to how she has been feeling in the previous 7 days.
2. All 10 items must be completed.
3. Care should be taken to avoid the possibility of the mother's discussing her answers with others.
4. The mother should complete the scale herself, unless she has limited English or has difficulty with reading.

Name: _____ Baby's age: _____
 Address: _____

Please **UNDERLINE** the answer that comes closest to how you have felt **IN THE PAST 7 DAYS**, not just how you feel today.

I have been able to laugh and see the funny side of things.

- 0 As much as I always could
- 1 Not quite so much now
- 2 Definitely not so much now
- 3 Not at all

I have looked forward with enjoyment to things.

- 0 As much as I ever did
- 1 Rather less than I used to
- 2 Definitely less than I used to
- 3 Hardly at all

*I have blamed myself unnecessarily when things went wrong.

- 3 Yes, most of the time
- 2 Yes, some of the time
- 1 Not very often
- 0 No, never

I have been anxious or worried for no good reason.

- 0 No, not at all
- 1 Hardly ever
- 2 Yes, sometimes
- 3 Yes, very often

*I have felt scared or panicky for no very good reason.

- 3 Yes, quite a lot
- 2 Yes, sometimes
- 1 No, not much
- 0 No, not at all

*Things have been getting on top of me.

- 3 Yes, most of the time I haven't been able to cope at all
- 2 Yes, sometimes I haven't been coping as well as usual
- 1 No, most of the time I have coped quite well
- 0 No, I have been coping as well as ever

*I have been so unhappy that I have had difficulty sleeping.

- 3 Yes, most of the time
- 2 Yes, sometimes
- 1 Not very often
- 0 No, not at all

*I have felt sad or miserable.

- 3 Yes, most of the time
- 2 Yes, quite often
- 1 Not very often
- 0 No, not at all

*I have been so unhappy that I have been crying.

- 3 Yes, most of the time
- 2 Yes, quite often
- 1 Only occasionally
- 0 No, never

*The thought of harming myself has occurred to me.

- 3 Yes, quite often
- 2 Sometimes
- 1 Hardly ever
- 0 Never

**Items are reverse scored (ie, 3, 2, 1, 0). Responses are scored 0, 1, 2, and 3 according to increased severity of symptoms. Total score is calculated by adding the scores for each of the 10 items. Scores > 12 identify patients at risk for postpartum depression.*

Figure reproduced from Cox et al.⁵ Users may reproduce the scale without further permission providing they respect copyright by quoting the names of the authors, the title, and the source of the paper in all reproduced copies.

rates of 17%. When patients were followed into the postpartum period, rates were found to have decreased to 13% by 6 to 8 weeks after the birth (level II evidence). Evans et al⁷ followed a group of women through pregnancy to the postpartum period and found that levels of depression during pregnancy were comparable to those reported during the postpartum period (level II evidence).

Gotlib and colleagues¹ were among the first to point out a continuum of depression through pregnancy and into the postpartum period (level II evidence). Despite their observation, research during the past 25 years has focused on postpartum

depression, rendering pregnancy-related depression a secondary issue. While the DSM-IV-TR recognizes postpartum-onset mood disorders, it does not make any reference to depression during pregnancy.

Only during the last 4 years have we seen depression during pregnancy resurface as an area of importance in the literature. This is due in part to the increase in referrals of depressed pregnant patients to reproductive psychiatry programs. This increase might reflect patients' increasing willingness to report symptoms to their primary caregivers or a heightened awareness among physicians of the onset

of illness during pregnancy. The fact that depression, if left undiagnosed, can persist throughout pregnancy into the postpartum period is gaining widespread recognition today (level II evidence).^{1,2,7,8}

Risk factors

Many studies have delineated the risk factors associated with depression during pregnancy; risk factors are summarized in **Table 1**. Personal and family history of depression are substantial biologic risk factors (level II evidence).^{11,12} Major psychosocial risk factors are history of childhood abuse, domestic violence or marital conflict, substance abuse or smoking, inadequate social support, single motherhood, lower educational levels, and unemployment (level II evidence).^{1,11,13-16}

Table 1. Summary of risk factors for developing depression during pregnancy

BIOLOGIC
History of mood and anxiety disorders
History of postpartum depression
History of premenstrual dysphoric disorder
Family history of psychiatric illness
PSYCHOSOCIAL
History of childhood abuse
Younger age
Unplanned pregnancy
Ambivalence or negative feelings about the pregnancy
Single motherhood
Greater number of children
Limited social support
Domestic violence or marital conflict
Low level of education and unemployment
Substance abuse and smoking

Once at-risk patients have been identified, they can be monitored throughout pregnancy for early signs of illness, whether it is new onset, exacerbation, or relapse of an existing depression.

Treatment

Interventions for depression during pregnancy most commonly include antidepressant medications; psychotherapy (individual or group); bright-light therapy; and very rarely, electroconvulsive therapy (ECT).

The two most commonly used psychotherapies are interpersonal therapy (IPT), which focuses on improving social interactions and coping with life transitions, and cognitive behavioural therapy (CBT), which aims to adjust patients' self-defeating thought patterns. Using IPT during pregnancy has been shown to improve mood substantially after 16 weeks (level I evidence).¹⁷ Unfortunately, there is little research on the effectiveness of other non-pharmacologic treatments for prenatal depression. Although the effectiveness of CBT for PPD has been demonstrated (level I evidence),¹⁸ to date there is no research on its use for depression during pregnancy. The effectiveness of group therapy for depression during pregnancy has not been reported, although antenatal group therapy has been shown to be effective in preventing PPD (level II evidence).¹⁹

Pharmacologic treatment. Selective serotonin reuptake inhibitors (SSRIs) and selective serotonin and norepinephrine reuptake inhibitors (SNRIs), are most commonly used today. The older antidepressants, like the tricyclic antidepressants, are no longer used routinely as first-line treatment for major depression. The SSRIs include fluoxetine (eg, Prozac), paroxetine (Paxil), sertraline (eg, Zoloft), fluvoxamine (eg, Luvox), and citalopram (Celexa). Venlafaxine is an example of a SNRI antidepressant. Bupropion (eg, Wellbutrin) and mirtazapine (Remeron) are among the newer dual-action antidepressants we are seeing in pregnant women. All SSRIs, as well as venlafaxine (Effexor), have been found to cross the placenta. Paroxetine and sertraline pass through the placenta more slowly than fluoxetine (level II evidence),²⁰ but this finding must be interpreted with caution in clinical practice. If a patient has responded best to fluoxetine, for example, it is recommended that she continue taking fluoxetine throughout pregnancy. Because every mother and baby metabolize medication differently, no universal statement about the choice of a particular medication during pregnancy can be made. **Table 2**²¹⁻³⁸ summarizes research findings on SSRI and SNRI exposure during pregnancy.

When a woman is treated with antidepressants during pregnancy, both mother's and doctor's concerns relate to the risk of teratogenesis, neonatal

Table 2. Summary of research findings on the safety of taking antidepressants during pregnancy: *There was no increased risk of teratogenicity with any of the medications.*

MEDICATION (NO. OF BABIES EXPOSED)	OBSTETRIC COMPLICATIONS AND NEONATAL WITHDRAWAL	LONG-TERM EFFECTS
Fluoxetine ²¹⁻²⁷ (1532)	Third trimester: premature delivery or perinatal complications; higher doses might lead to low birth weight; transient withdrawal symptoms*	No long-term adverse effects; children of depressed mothers did not have cognitive and language difficulties
Sertraline ^{24,25,28-30} (234)	Transient withdrawal symptoms*	Further research required
Paroxetine ^{24,25,28,30,31} (313)	Transient withdrawal symptoms*	Further research required
Citalopram ^{22-24,30,32,33} (318) Swedish birth registry (375); Lundbeck Safety Database (100)	None	Further research required
Fluvoxamine ²⁸⁻³⁰ (30)	None	Further research required
Venlafaxine ^{24,29,34} (174)	Transient withdrawal symptoms*	Further research required
Mirtazapine ^{35,36} (9)	None	Further research required
Bupropion ³⁷ (534)	None	Further research required
Trazodone ³⁸ (58) Nefazodone ³⁸ (89)	None	Further research required

*Transient neonatal withdrawal symptoms include respiratory distress, jaundice, jitteriness, increased fussing, tremors, and increased crying.

toxicity, and long-term effects on child development.³⁹ During the first trimester, the main concern is malformation of the fetus, although there is no current evidence that SSRIs or venlafaxine cause increased teratogenicity (level I evidence).^{28,40,41} Concerns in the third trimester focus on neonatal withdrawal because third-trimester exposure to antidepressants has been correlated with higher risk of adverse effects in neonates, such as respiratory distress, feeding difficulties, and low birth weight. These effects, however, have been shown to be transient (level I evidence).²³⁻²⁵ Although there is little evidence from research, no long-term adverse effects from prenatal exposure to SSRIs or venlafaxine have been reported (level II evidence).^{26,27,42} This area requires ongoing research.

A mother's mood should be monitored very closely throughout pregnancy, particularly in the third trimester, where somatic changes might lead to fluctuating dose requirements (level II evidence).^{43,44} In a recent study, pharmacologically undertreated maternal mood disorders during the third trimester were found to contribute to poor neonatal adaptation (level II evidence).⁴⁵

Health Canada recently issued a caution about use of SSRIs during the third trimester. Although mothers and doctors might be concerned about exposure, rapid discontinuation of medications is not recommended because it can substantially increase risk of relapse (level II evidence).⁴⁶ It is always important to inform obstetricians and neonatologists about exposure to SSRIs in utero so that babies can be closely monitored during delivery and receive appropriate treatment, if necessary.

Among the more than 400 case reports on short- and long-term effects of exposure to tricyclic medications during pregnancy, few major adverse effects have been reported.^{47,48} In spite of this, because of their side effect profile, tricyclics are used less frequently today. Monoamine oxidase inhibitors are not recommended during pregnancy.

Nonpharmacologic biologic treatments. Bright-light therapy could be an alternative to pharmacologic intervention and has been shown to be effective for treating antenatal depression (level II and III evidence).^{49,50} It is particularly helpful for patients who report seasonal variation in their moods and who do not want to use antidepressants during pregnancy.

Several case reports have shown ECT to be relatively safe and effective during pregnancy (level II evidence).⁵¹⁻⁵³ The American Psychiatric Association, however, has strict guidelines for administration of ECT⁵⁴: it should be used only when women are severely psychotic or acutely suicidal and when other therapies or medications have clearly failed.

Treatment of depression during pregnancy requires patients and physicians to make collaborative decisions. Each treatment needs to be chosen on a case-by-case basis. For example, women with

a history of recurrent depressive episodes, who are already taking medication at conception, should be encouraged to continue the medication through pregnancy to the postpartum period. If a patient has a history of severe depression and experiences a relapse during her current pregnancy, recommencing treatment with an antidepressant to which she has responded favourably in the past is recommended. For patients who are experiencing an initial depressive episode during pregnancy, treatment with an antidepressant is recommended only if the depression is severe and unlikely to respond to psychotherapy.

The benefits of treating depression during pregnancy far outweigh the consequences that can result from undiagnosed and untreated depression. Maternal psychiatric illness is associated with poor prenatal behaviour, including low attendance at prenatal checkups and increased substance use (level II evidence).^{55,56} Adverse obstetric and neonatal outcomes, such as increased preterm delivery, low birth weight, and admission to neonatal nurseries, have also been linked to depression during pregnancy (level II evidence).⁵⁷⁻⁶⁰ If depression persists into the postpartum period, it can have long-term consequences for both mother and baby. Mothers might go on to develop chronic mood disorders, and untreated postpartum depression can impair mother-infant attachment (level I evidence).^{61,62} Finally, being exposed to a chronically depressed mother can have cognitive, emotional, and behavioural consequences for a child (level II evidence).⁶³⁻⁶⁵

Conclusion

Early detection and treatment of antenatal depression is vital. Once diagnosis is confirmed, family physicians need to include patients in a risk-benefit analysis and carefully outline the consequences of untreated depression for both mother and baby. For mild-to-moderate depression, nonpharmacologic treatments should be offered first, along with referral to a psychologist, if available. For more seriously ill patients, psychotherapy alone might be insufficient, and additional treatment with an antidepressant might be required.

Family physicians might not feel comfortable treating seriously depressed women during pregnancy. Mothers themselves might want a specialist involved in the hope of avoiding taking medications during pregnancy or of learning more about the safety of using medications during pregnancy. Referral to a psychiatrist for expert opinion and ongoing treatment should be considered.

Most research to date supports favourable outcomes for women and babies exposed to antidepressants during pregnancy (level II and III evidence).⁶⁶ It is clear now that comprehensive treatment plans must include not only patients and their doctors, but also where applicable, women's partners, obstetricians, neonatologists, and other caregivers (eg, midwives, counselors). Once treatment has been started, it is critical to monitor and follow up into the postpartum period to prevent relapse. In the last 25 years, we have learned much about postpartum depression. We hope the gains made in this area will help us better understand depression during pregnancy.



Competing interests

None declared

Correspondence to: Deirdre Ryan, Reproductive Mental Health Program, BC Women's Hospital, H214-4500 Oak St, Vancouver, BC V6H 3N1; telephone (604) 875-2025; fax (604) 875-3136; e-mail dryan@cw.bc.ca

References

1. Gotlib IH, Whiffen VE, Mount JH, Milne K, Cordy NI. Prevalence rates and demographic characteristics associated with depression in pregnancy and the postpartum. *J Consult Clin Psychol* 1989;57(2):269-74.
2. Austin MP. Antenatal screening and early intervention for "perinatal" distress, depression and anxiety: where to from here? *Arch Women Ment Health* 2004;7(1):1-6.
3. Chaudron LH, Klein MH, Remington P, Palta M, Allen C, Essex MJ. Predictors, prodromes and incidence of postpartum depression. *J Psychosom Obstet Gynaecol* 2001;22(2):103-12.
4. American Psychiatric Association. *Diagnostic and statistical manual of mental disorders*. 4th ed, text rev. Washington, DC: American Psychiatric Association; 2000.
5. Cox JL, Holden JM, Sagovsky R. Detection of postnatal depression. Development of the 10-item Edinburgh Postnatal Depression Scale. *Br J Psychiatry* 1987;150:782-6.
6. Murray D, Cox J. Screening for depression during pregnancy with the Edinburgh Postnatal Depression Scale (EPDS). *J Reprod Infant Psychol* 1990;8:99-107.
7. Evans J, Heron J, Francomb H, Oke S, Golding J. Cohort study of depressed mood during pregnancy and after childbirth. *BMJ* 2001;323(7307):257-60.
8. Josefsson A, Berg G, Nordin C, Sydsjö G. Prevalence of depressive symptoms in late pregnancy and postpartum. *Acta Obstet Gynecol Scand* 2001;80(3):251-5.
9. Bennett HA, Einarson A, Taddio A, Koren G, Einarson TR. Prevalence of depression during pregnancy: systematic review. *Obstet Gynecol* 2004;103(4):698-709.
10. Beck AT, Ward CH, Mendelson M, Mock J, Erbaugh J. An inventory for measuring depression. *Arch Gen Psychiatry* 1961;4:561-71.
11. Marcus SM, Flynn HA, Blow FC, Barry KL. Depressive symptoms among pregnant women screened in obstetrics settings. *J Womens Health (Larchmt)* 2003;12(4):373-80.
12. Altshuler LL, Hendrick V, Cohen LS. Course of mood and anxiety disorders during pregnancy and the postpartum period. *J Clin Psychiatry* 1998;59(Suppl 2):29-33.
13. Farber EW, Herbert SE, Reviere SL. Childhood abuse and suicidality in obstetrics patients in a hospital-based urban prenatal clinic. *Gen Hosp Psychiatry* 1996;18(1):56-60.

14. Pajulo M, Savonlahti E, Sourander A, Helenius H, Piha J. Antenatal depression, substance dependency and social support. *J Affect Disord* 2001;65(1):9-17.
15. O'Hara MW. Social support, life events, and depression during pregnancy and the puerperium. *Arch Gen Psychiatry* 1986;43:569-73.
16. Bolton HL, Hughes PM, Turton P, Sedgwick P. Incidence and demographic correlates of depressive symptoms during pregnancy in an inner London population. *J Psychosom Obstet Gynaecol* 1998;19(4):202-9.
17. Spinelli MG, Endicott J. Controlled clinical trial of interpersonal psychotherapy versus parenting education program for depressed pregnant women. *Am J Psychiatry* 2003;160(3):555-62.
18. Misri S, Reebye P, Corral M, Milis L. The use of paroxetine and cognitive-behavioral therapy in postpartum depression and anxiety: a randomized controlled trial. *J Clin Psychiatry* 2004;65:1236-41.
19. Zlotnick C, Johnson SL, Miller IW, Pearlstein T, Howard M. Postpartum depression in women receiving public assistance: pilot study of an interpersonal-therapy-oriented group intervention. *Am J Psychiatry* 2001;158(4):638-40.
20. Hendrick V, Stowe ZN, Altschuler LL, Hwang S, Lee E, Haynes D. Placental passage of antidepressant medications. *Am J Psychiatry* 2003;160(5):993-6.
21. Heikkinen T, Ekblad U, Palo P, Laine K. Pharmacokinetics of fluoxetine and norfluoxetine in pregnancy and lactation. *Clin Pharmacol Ther* 2003;73(4):330-7.
22. Laine K, Heikkinen T, Ekblad U, Kero P. Effects of exposure to selective serotonin reuptake inhibitors during pregnancy on serotonergic symptoms in newborns and cord blood monoamine and prolactin concentrations. *Arch Gen Psychiatry* 2003;60:720-6.
23. Zeskind PS, Stephens LE. Maternal selective serotonin reuptake inhibitor use during pregnancy and newborn neurobehavior. *Pediatrics* 2004;113(2):368-75.
24. Kallen B. Neonate characteristics after maternal use of antidepressants in late pregnancy. *Arch Pediatr Adolesc Med* 2004;158(4):312-6.
25. Oberlander TF, Misri S, Fitzgerald CE, Kostaras X, Rurak D, Riggs W. Pharmacologic factors associated with transient neonatal symptoms following prenatal psychotropic medication exposure. *J Clin Psychiatry* 2004;65(2):230-7.
26. Nulman I, Rovet J, Stewart DE, Wolpin J, Gardner HA, Theis JG, et al. Neurodevelopment of children exposed in utero to antidepressant drugs. *N Engl J Med* 1997;336(4):258-62.
27. Nulman I, Rovet J, Stewart DE, Wolpin J, Pace-Asciak P, Shuhaiber S, et al. Child development following exposure to tricyclic antidepressants or fluoxetine throughout fetal life: a prospective, controlled study. *Am J Psychiatry* 2002;159(11):1889-95.
28. Kuln NA, Pastuszak A, Sage SR, Schick-Boschetto B, Spivey G, Feldkamp M, et al. Pregnancy outcome following maternal use of the new selective serotonin reuptake inhibitors: a prospective controlled multicenter study. *JAMA* 1998;279(8):609-10.
29. Hostetter A, Ritchie JC, Stowe ZN. Amniotic fluid and umbilical cord blood concentrations of antidepressants in three women. *Biol Psychiatry* 2000;48(10):1032-4.
30. Hendrick V, Smith LM, Suri R, Hwang S, Haynes D, Altschuler L. Birth outcomes after prenatal exposure to antidepressant medication. *Am J Obstet Gynecol* 2003;188(3):812-5.
31. Costei AM, Kozar E, Ho T, Ito S, Koren G. Perinatal outcome following third trimester exposure to paroxetine. *Arch Pediatr Adolesc Med* 2002;156(11):1129-32.
32. Heikkinen T, Ekblad U, Kero P, Ekblad S, Laine K. Citalopram in pregnancy and lactation. *Clin Pharmacol Ther* 2002;72(2):184-91.
33. Ericson A, Kallen B, Wiholm B. Delivery outcome after the use of antidepressants in early pregnancy. *Eur J Clin Pharmacol* 1999;55(7):503-8.
34. Einarson A, Fatoye B, Sarkar M, Lavigne SV, Brochu J, Chambers C, et al. Pregnancy outcome following gestational exposure to venlafaxine: a multicenter prospective controlled study. *Am J Psychiatry* 2001;158(10):1728-30.
35. Kesim M, Yaris F, Kadioglu M, Yaris E, Kalyoncu NI, Ulku C. Mirtazapine use in two pregnant women: is it safe? *Teratology* 2002;66:204.
36. Saks BR. Mirtazapine: treatment of depression, anxiety, and hyperemesis gravidarum in the pregnant patient. A report of 7 cases. *Arch Women Ment Health* 2003;3:165-70.
37. Bupropion Pregnancy Registry. *Interim report 9/1/97-2/29/04*. Research Triangle Park, NC: Glaxo SmithKline; 2004. p. 1-29.
38. Einarson A, Bonari L, Voyer-Lavigne S, Addis A, Matsui D, Johnson Y, et al. A multicenter prospective controlled study to determine the safety of trazodone and nefazodone use during pregnancy. *Can J Psychiatry* 2003;48(2):106-10.
39. Cohen LS, Nonacs R, Viguera AC, Remnick A. Diagnosis and treatment of depression during pregnancy. *CNS Spectr* 2004;9(3):209-16.
40. Simon GE, Cunningham ML, Davis RL. Outcomes of prenatal antidepressant exposure. *Am J Psychiatry* 2002;159(12):2055-61.
41. McElhatton PR, Garbis HM, Elefant E, Vial T, Bellemine B, Mastroiacovo P, et al. The outcome of pregnancy in 689 women exposed to therapeutic doses of antidepressants. A collaborative study of the European Network of Teratology Information Services (ENTIS). *Reprod Toxicol* 1996;10(4):285-94.
42. Loebstein R, Koren G. Pregnancy outcome and neurodevelopment of children exposed in utero to psychoactive drugs: the Motherisk experience. *J Psychiatry Neurosci* 1997;22(3):192-6.
43. Altschuler LL, Hendrick VC. Pregnancy and psychotropic medication: changes in blood levels. *J Clin Psychopharmacol* 1996;16(1):78-80.
44. Hostetter A, Stowe ZN, Strader JR Jr, McLaughlin E, Llewellyn A. Dose of selective serotonin uptake inhibitors across pregnancy: clinical implications. *Depress Anxiety* 2000;11(2):51-7.
45. Misri S, Oberlander TF, Fairbrother N, Carter D, Ryan D, Kuan AJ, et al. Relation between prenatal maternal mood and anxiety and neonatal health. *Can J Psychiatry* 2004;49(10):684-9.
46. Einarson A, Selby P, Koren G. Abrupt discontinuation of psychotropic drugs during pregnancy: fear of teratogenic risk and impact of counselling. *J Psychiatry Neurosci* 2001;26(1):44-8.
47. Altschuler LL, Cohen L, Szuba MP, Burt VK, Gitlin M, Mintz J. Pharmacologic management of psychiatric illness during pregnancy: dilemmas and guidelines. *Am J Psychiatry* 1996;153(5):592-606.
48. Cohen LS, Rosenbaum JF. Psychotropic drug use during pregnancy: weighing the risks. *J Clin Psychiatry* 1998;59(Suppl 2):18-28.
49. Oren DA, Wisner KL, Spinelli M, Epperson CN, Peindl KS, Terman JS, et al. An open trial of morning light therapy for treatment of antepartum depression. *Am J Psychiatry* 2002;159(4):666-9.

EDITOR'S KEY POINTS

- Pregnancy is a high-risk time for new-onset or reactivated depression; if untreated, depression can continue into the postpartum period.
- A history of depression, especially postpartum depression, is a risk factor for reactivation of mood disorders during pregnancy and afterward.
- For mild-to-moderate depression, psychotherapy might be sufficient. For severe illness, antidepressant medications are recommended and safe. Medication decisions should be made on a case-by-case basis in consultation with each patient.
- Neonatal withdrawal effects are transient, and the effects of untreated depression are of more concern than neonatal exposure to the medication.

POINTS DE REPÈRE DU RÉDACTEUR

- Durant la grossesse, le risque de développer ou de réactiver une dépression est accru; sans traitement, la dépression peut se prolonger durant le postpartum.
- Une dépression antérieure, notamment durant le postpartum, augmente le risque d'une réactivation des troubles de l'humeur durant la grossesse.
- Dans les cas légers à modérés, la psychothérapie pourrait suffire. Pour une dépression sévère, l'usage d'antidépresseurs est recommandé et sécuritaire. L'option pharmacologique doit être individualisée et discutée avec la patiente.
- Les effets du sevrage néonatal sont transitoires; ceux d'une dépression non traitée sont plus préoccupants que l'exposition du nouveau-né à la médication.

50. Epperson CN, Terman M, Terman JS, Hanusa BH, Oren DA, Peindl KS, et al. Randomized clinical trial of bright light therapy for antepartum depression: preliminary findings. *J Clin Psychiatry* 2004;65(3):421-5.
51. Bhatia SC, Baldwin SA, Bhatia SK. Electroconvulsive therapy during the third trimester of pregnancy. *J ECT* 1999;15(4):270-4.
52. Miller LJ. Use of electroconvulsive therapy during pregnancy. *Hosp Community Psychiatry* 1994;45(5):444-50.
53. Polster DS, Wisner KL. ECT-induced premature labor: a case report. *J Clin Psychiatry* 1999;60(1):53-4.
54. Weiner RD, Coffey CE, Fochtmann L, Greenberg R, Isenberg KE, Moench L, et al, and American Psychiatric Association. *The practice of ECT: recommendations for treatment, training and privileging*. 2nd ed. Washington, DC: American Psychiatric Press; 2001.
55. Lindgren K. Relationships among maternal-fetal attachment, prenatal depression, and health practices in pregnancy. *Res Nurs Health* 2001;24(3):203-17.
56. Zuckerman B, Amaro H, Bauchner H, Cabral H. Depressive symptoms during pregnancy: relationship to poor health behaviors. *Am J Obstet Gynecol* 1989;160(5 Pt 1):1107-11.
57. Chung TK, Lau TK, Yip AS, Chiu HF, Lee DT. Antepartum depressive symptomatology is associated with adverse obstetric and neonatal outcomes. *Psychosom Med* 2001;63(5):830-4.
58. Hedegaard M, Henriksen TB, Sabroe S, Secher NJ. Psychological distress in pregnancy and preterm delivery. *BMJ* 1993;307(6898):234-9.
59. Dayan J, Creveuil C, Herlicovitz M, Herbel C, Baranger E, Savoye C, et al. Role of anxiety and depression in the onset of spontaneous preterm labor. *Am J Epidemiol* 2002;155(4):293-301.
60. Kurki T, Hilesmaa V, Raitasalo R, Mattila H, Ylikorkala O. Depression and anxiety in early pregnancy and risk for preeclampsia. *Obstet Gynecol* 2000;95(4):487-90.
61. Beck CT. The effects of postpartum depression on maternal-infant interaction: a meta-analysis. *Nurs Res* 1995;44(5):298-304.
62. Edhborg M, Lundh W, Seimyr L, Widstrom AM. The parent-child relationship in the context of maternal depressive mood. *Arch Women Ment Health* 2003;6(3):211-6.
63. Righetti-Veltama M, Bousquet A, Manzano J. Impact of postpartum depressive symptoms on mother and her 18-month-old infant. *Eur Child Adolesc Psychiatry* 2003;12(2):75-83.
64. Hay DF, Pawlby S, Angold A, Harold GT, Sharp D. Pathways to violence in the children of mothers who were depressed postpartum. *Dev Psychol* 2003;39(6):1083-94.
65. Righetti-Veltama M, Conne-Perreard E, Bousquet A, Manzano J. Postpartum depression and mother-infant relationship at 3 months old. *J Affect Disord* 2002;70(3):291-306.
66. Wisner KL, Zarin DA, Holmboe ES, Appelbaum PS, Gelenberg AJ, Leonard HL, et al. Risk-benefit decision making for treatment of depression during pregnancy. *Am J Psychiatry* 2000;157(12):1933-40.